

Coppertone UltraGuard SPF 70 Sunscreen Lotion

Drug Facts	
Active ingredients	Purpose
Avobenzone 3%, Homosalate 15%, Octisalate 5%, Octocrylene 10%, Oxybenzone 6%.....Sunscreen	
Uses	
<ul style="list-style-type: none"> • helps prevent sunburn • If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun 	
Warnings	
For external use only	
Do not use on damaged or broken skin	
When using this product keep out of eyes. Rinse with water to remove	
Stop use and ask a doctor if rash occurs	
Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> • apply liberally 15 minutes before sun exposure • reapply: <ul style="list-style-type: none"> • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: <ul style="list-style-type: none"> • limit time in the sun, especially from 10 a.m.– 2 p.m. • wear long-sleeve shirts, pants, hats, and sunglasses • children under 6 months: Ask a doctor 	
Other information	
<ul style="list-style-type: none"> • protect this product from excessive heat and direct sun • may stain or damage some fabrics or surfaces 	
Inactive ingredients	
Water, Butylene Glycol, Microcrystalline Cellulose, Glyceryl Stearate, Behenyl Alcohol, Benzyl Alcohol, Diethylhexyl Syringylidenemalonate, Tocopherol (Vitamin E), Retinyl Palmitate (Vitamin A), Sodium Ascorbyl Phosphate, Stearic Acid, Palmitic Acid, Lauryl Alcohol, Myristyl Alcohol, Cetyl Alcohol, Lecithin, Caprylic/Capric Triglyceride, Chlorphenesin, Cellulose Gum, Butylated PVP, Disodium EDTA	
Questions? 866-288-3330	